

The vascular plants in the following two releves are part of a series of 100 releves prepared by Hans Roemer in 1996 to describe and classify “pocket grasslands” found on SE Vancouver Island (transcribed 2011, names updated to present synonyms).

Locality: Grass balds W of Veitch Creek (approximate UTM 10U 456110, 5363240)

Cover values are from 1 (10%) to 10 (100% cover); < = less than 1%.

Releve # >>	3378	3379
<i>Festuca roemerii</i>	6	7
<i>Anthoxanthum odoratum</i>	1	3
<i>Achillea millefolium</i>	2	1
<i>Danthonia californica</i>	2	1
<i>Camassia quamash</i>	1	1
<i>Lomatium utriculatum</i>	1	1
<i>Plectritis congesta</i>	1	1
<i>Hypochaeris radicata</i>	<	2
<i>Aira caryophylla</i>	<	1
<i>Montia parvifolia</i>	<	1
<i>Agrostis cf. pallens</i>	2	
<i>Aira praecox</i>	<	<
<i>Bromus hordeaceus</i>	2	
<i>Carex inops</i>	3	
<i>Phlox gracilis</i>	<	<
<i>Rumex acetosella</i>	<	<
<i>Brodiaea coronaria</i>	1	
<i>Cerastium arvense</i>		1
<i>Eriophyllum lanatum</i>	1	
<i>Lotus micranthus</i>	1	
<i>Luzula multiflora</i>		1
<i>Mahonia aquifolium</i>	1	
<i>Allium acuminatum</i>	<	
<i>Clarkia amoena</i>	<	
<i>Collinsia parviflora</i>		<
<i>Epilobium minutum</i>	<	
<i>Galium aparine</i>		<
<i>Olsynium douglasii</i>		<
<i>Trifolium tridentatum</i>		<